

# Victor® Moleworms™ Kills Moles

## How to Apply: Treating Subsurface Runways

### Locate active tunnels



Make a 1 inch hole in the roof of a mole surface tunnel. Do not crush tunnel. Always mark open tunnels. Mark these openings to allow you to revisit later.



Revisit tunnel openings 24-48 hours later. Tunnels that have had holes resealed within 48 hours are an indication of mole activity and a good place to bait.

### Place the product in tunnels




Poke a new hole in the roof of the active tunnel just larger than the diameter of the product.



Drop the product down into the hole and reseat with dirt. Repeat the process for every 5 to 10 feet of active tunnel.

### Checking the results

Five days after placing the product into the active tunnels, use procedure above to see if tunnel remains active and if so, retreat any tunnels that still appear to be active.



EPA Reg. No. 349-19  
EPA Est. No. 47629-PA-001

Model No. M6009PCO  
Manufactured for:  
Senorel Chemical Company, Inc.  
89 N. Locust St  
Lititz, PA 17543  
855-5-VICTOR



# Victor® Moleworms™ Kills Moles

## 25 Worms

10 Active Tunnel  
Locator Flags

3 Protective  
Gloves

Moleworms™ resemble real earthworms!  
Moles eat earthworms as part of their diet.



This Product May Only be Used to Control Moles in Manual, Below-Ground Applications

KEEP OUT OF REACH  
OF CHILDREN  
CAUTION

Señalar Prohibido  
PARA LOS NIÑOS  
PRECAUCIÓN

MANTÉNGASE FUERA DEL  
ALCANCE DE LOS NIÑOS  
PRECAUCIÓN

No es permitido para niños  
MANTÉNGASE FUERA DEL  
ALCANCE DE LOS NIÑOS

ACTIVE INGREDIENT PRECAUTIONARY:  
Bromethalin 0.025%  
OTHER INGREDIENTS PRECAUTIONARY:  
Talc 99.975%  
TOTAL 100.000%

NET WEIGHT: 5.75 OZ (162.5g)

### DIRECTIONS FOR USE:

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

**READ THIS LABEL:** Read this entire label and follow all use directions and use precautions.

**Important:** Do not expose children to this product or use it for any purpose other than to control moles. To help prevent accidents:

1) Store product not in use in locations out of the reach of children and pets. 2) Follow all application directions and USE RESTRICTIONS on this label. Apply bait only in underground runways of moles. Keep pets out of treated areas. Do NOT use this product above ground. 3) Dispose of product container and unused or spilled product as specified on label.

**Use restrictions:** This product may only be applied at the use sites and by the application methods indicated on this label.

This product may only be used in lawns, parks, around homes, ornamental gardens, nurseries and other non-crop grassy areas to control eastern moles (*Scalopus aquaticus*, star-nosed moles (*Geopline arifolia*) or Skinkapops spp. Bait must be applied directly into moles' burrow systems. This bait must be applied underground in either main underground runways or in subsurface feeding tunnels. This bait may not be used above ground. Wear gloves when bait is handled.

**How to select proper treatment areas:** The presence of moles is indicated by a network of subsurface runways in the turf caused by moles foraging for food just below the soil surface.

You may also see a series of conical mounds of earth pushed up from deep burrows.

### Treating conical mounds

Moles construct deeper tunnels that are not visible from the surface. These deep tunnel systems are often marked by a series of conical mounds of loose dirt 4 to 12 inches high formed when dirt is pushed to the surface through a mole tunnel shaft.

1. Remove all of the rounded loose dirt that is at ground level.
2. Continue to remove loose dirt that is below ground level and you will eventually locate the primary tunnel shaft.
3. Once the main tunnel shaft has been discovered and opened, leave it opened and return in 48 hours to see if the opening has been resealed with loose dirt. This indicates that this mound and tunnel shaft is in use by a mole.
4. Reopen the tunnel shaft and drop the product down into the shaft and reseat opening with dirt.
5. Revisit this mound in 5 days and check again to see if it remains active.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**Pesticide Storage:** Store only in original container, in a dry place inaccessible to children and pets. Keep containers closed and away from other chemicals.

**Pesticide Disposal and Container Handling:** Nonrefillable Container. Do not reuse or refill this container.

If empty: Place in trash or other for recycling if available.

If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any floor or outdoor drain.

### PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed. Keep away from children, domestic animals and pets. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

**First Aid**

Have the product container or label with you when calling a poison control center or going for treatment.

If swallowed:

- Call a poison control center, doctor, or 1-800-222-1222 (nights, weekends) or 1-800-837-7544 (Monday - Friday, daytime) immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting, unless told to do so by the poison control center or doctor.

**Treatment for pet poisoning:** If an animal eats bait, call veterinarian or numbers listed above at once.

**Not to Physicians or Veterinarians:**

Contains the nerve poison Bromethalin.

This is not an anticoagulant type rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of central adrena and should be treated accordingly through administration of an osmotic diuretic and corticosteroids.

### Environmental Hazards:

This product is extremely toxic to mammals and birds. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not contaminate water when disposing of equipment wash water or residue. Do not apply directly to water.



DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READ THIS LABEL: Read this entire label and follow all use directions and use precautions. See outer container for complete directions.

Important: Do not expose children to this product or use it for any purpose other than to control moles. To help prevent accidents: 1) Store product not in use in locations out of the reach of children and pets. 2) Follow all application directions and **USE RESTRICTIONS** on this label. Apply bait only in underground runways of moles. Keep pets out of treated areas. Do NOT use this product above ground. 3) Dispose of product container and unused or spoiled product as specified on label.

Use restrictions: This product may only be applied at the use sites and by the application methods indicated on this label. This product may only be used in lawns, parks, around homes, ornamental gardens, nurseries and other non-crop grassy areas to control eastern moles (*Scalopus aquaticus*), star-nosed moles (*Corylura cristata*) or *Scapanus* spp. Bait must be applied directly into moles' burrow systems. This bait must be applied

underground in either main underground runways or in subsurface feeding tunnels. This bait may not be used above ground. Wear gloves when bait is handled.

How to select proper treatment areas: The presence of moles is indicated by a network of subsurface runways in the turf caused by moles foraging for food just below the soil surface. You may also see a series of conical mounds of earth pushed up from deep burrows.

Treating conical mounds

Moles construct deeper tunnels that are not visible from the surface. These deep tunnels systems are often marked by a series of conical mounds of loose dirt 4 to 12 inches high formed when dirt is pushed to the surface through a mole tunnel shaft. 1) Remove all of the mounded loose dirt that is at ground level. 2) Continue to remove loose dirt that is below ground level and you will eventually locate the primary tunnel shaft. 3) Once the main tunnel shaft has been discovered and opened, leave it opened and return in 48 hours to see if the opening has been reclosed with loose dirt. This indicates that this mound and tunnel shaft is in use by a mole. 4) Reopen the tunnel shaft and drop the product down into the shaft and reclose opening with dirt. 5) Revisit this mound in 5 days and check again to see if it remains active.

Victor<sup>®</sup>  
Moleworms<sup>™</sup>  
Kills Moles

This Product May Only be Used to Control Moles in Manual, Below-Ground Applications

5 Worms

NET WEIGHT: 1.15 OZ. (32.5g)

KEEP OUT OF REACH  
OF CHILDREN  
CAUTION

See Rear Panel for  
FIRST AID and other  
PRECAUTIONARY STATEMENTS

ACTIVE INGREDIENT:	
Bromethalin	0.025%
OTHER INGREDIENTS:	99.975%
TOTAL	100.000%

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **PESTICIDE STORAGE:** Store only in original container in a dry place inaccessible to children and pets. Keep containers closed and away from other chemicals. **PESTICIDE DISPOSAL AND CONTAINER HANDLING:** Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

**CAUTION:** Harmful if swallowed. Keep away from children, domestic animals and pets. Any person who retrieves carcasses or unused bait following application of this product must wear gloves.

Manufactured for:  
Senoret Chemical Company, Inc.

69 N. Locust St  
Lititz, PA 17543  
855-5-VICTOR

First Aid

Have the product container or label with you when calling a poison control center or going for treatment.

If swallowed:

Call a poison control center, doctor, or 1-800-222-1222 (nights, weekends) or 1-800-837-7644 (Monday – Friday, daytime) immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor.

**Treatment for pet poisoning:** If an animal eats bait, call veterinarian or numbers listed above at once.

**Note to Physician or Veterinarian:** Contains the nerve poison Bromethalin. This is not an anticoagulant type rodenticide. For humans or animals ingesting bait and/or showing poisoning signs (such as muscle tremors, loss of hind limb use, or seizures for animals), limit absorption by either emesis or gastric lavage. Sublethal symptoms, if present, would be the result of cerebral edema and should be treated accordingly through administration of an osmotic diuretic and corticosteroid.

Environmental Hazards

This product is extremely toxic to mammals and birds. Dogs and other predatory and scavenging mammals and birds might be poisoned if they feed upon animals that have eaten this bait. Do not contaminate water when disposing of equipment wash water or rinsate. Do not apply directly to water.